Chief Medical Officer, Public Health and Sport Directorate

T: 0131-244 2164 F: 0131-244 2157
E: joyce.whytock@scotland.gsi.gov.uk

The Scottish Government

CARBON MONOXIDE (CO) POISONING: NEEDLESS DEATHS, UNNECESSARY INJURY

Dear Colleague

The Health Protection Agency (HPA) and the Health and Safety Executive (HSE) will once again be supporting the UK wide Carbon Monoxide (CO) awareness week, organised by the charity CO Awareness, commencing Monday 21 November 2011 - **links provided at the end of Annex A.**

CO poisoning should be a problem of the past, but unfortunately, this is not so. A number of people are still dying from accidental acute CO poisoning and many more than previously realised are injured from sub-lethal poisonings, which can often lead to lasting neurological damage in victims. These victims are exposed to CO in their homes, but are also at risk from exposure in holiday residences, including caravans.

There are approximately 2 accidental deaths from acute CO poisoning in Scotland every year and there were around 135 patients who had a diagnosis of 'Toxic Effect of Carbon Monoxide' recorded at a hospital episode between 1 January 2008 and 31 December 2010. Of these patients, approximately 115 patients had 'Toxic Effect of Carbon Monoxide' recorded as a main diagnosis. Unfortunately, this does not provide any real indication of the severity of a particular case.

However, there is new data which suggests that there are likely to be even more cases of non-fatal poisonings in people who attend A&E, are treated for CO poisoning, but who do not require admission to hospital – this is of great concern as CO poisoning can lead to chronic health problems. The number of people exposed to CO, but who are unaware of the cause and do not present at their GPs surgery or local hospital is still not known but is likely to be many more.

Yet these deaths and accidental poisonings can be prevented: through greater public awareness and in particular, increased vigilance amongst health professionals of the signs and symptoms of exposure in their patients.

From the Chief Medical Officer
Chief Nursing Officer
Sir Harry Burns MPH FRCS(Glas) FRCP(Ed) FFPH
Ros Moore RGN, RNT, BSc (Hons) Nursing, MA

17 November 2011

SGHD/CMO(2011)14

For action
Chief Executives, NHS Boards; Nurse Directors, NHS Boards; General Practitioners; Practice Managers; Practice Nurses; Health Visitors; CPHMs; Directors of Public Health; Medical Directors, NHS Boards; Health Protection Teams; Scottish Ambulance Service; NHS 24; Chief Executives, Local Authorities; Directors of Education, Local Authorities; Universities Scotland; Learning and Teaching Scotland

For information
Chairs, NHS Boards;
Chairs, Local Authorities;
Consultant Paediatricians;
Consultant Physicians;
Directors of Pharmacy;
Health Protection Scotland;
Chief Executive, NHS Health Scotland

Further Enquiries
Policy issues
Joyce Whytock
3ES, St. Andrews House
joyce.whytock@scotland.gsi.gov.uk

Scientific issues
Dr Arthur Johnston
2N.07, St. Andrews House
arthur.johnston@scotland.gsi.gov.uk

Dr Colin Ramsay
Health Protection Scotland
colin.ramsay@nhs.net

INVESTOR IN PEOPLE
POSITIVE ABOUT DISABLED PEOPLE

An information leaflet for patients and the public is also available (links provided at the **end of Annex A**) giving guidance on preventing CO poisoning and what to do if they suspect they have been poisoned and to enable you to correctly advise your patient.

Unfortunately, on occasions in the past, colleagues have advised patients to stay at home and keep warm when the symptoms they presented with were in fact those of CO poisoning – the result can be fatal.

I hope you find the information helpful and thank you in advance in ensuring a successful UK wide CO awareness campaign during 2011.

Yours sincerely

Harry Burns *Ros Moore*

HARRY BURNS **ROS MOORE**

INVESTOR IN PEOPLE

Summary
This letter brings together the most up-to-date information on carbon monoxide (CO) poisoning. It describes the signs and symptoms which should be looked for; explains the investigations that may be necessary to establish whether CO poisoning has occurred; describes how cases should be managed; sets out the main sources of CO in the home and provides sources of further advice and information.

Background
Carbon monoxide (CO) is a colourless, odourless gas that causes the accidental deaths of approximately 2 people each year in Scotland. Furthermore, around 115 people each year are sufficiently ill to need hospital treatment and new data suggests that a similar number of people are treated but not admitted to hospital each year from accidental poisoning by CO. Poisoning by CO is almost certainly under-diagnosed and there could well be a large number of people being exposed and suffering the ill effects of exposure. Older people, children, pregnant women and their babies and those with breathing problems or cardiovascular disease are at increased risk. Poisoning can result in lasting neurological damage.

How to diagnose carbon monoxide poisoning
Recognising CO poisoning is not at all easy, as it may simulate many other conditions and unless poisoning is suspected, the diagnosis will be missed.

The onset of symptoms is often insidious and may not be recognised by either the patient or the doctor. The commonest symptoms and signs and an indication of their approximate frequency in CO poisoning are shown below:

Headache	90% of cases
Nausea and vomiting	50% of cases
Vertigo	50% of cases
Alteration in consciousness	30% of cases
Subjective weakness	20% of cases

While chronic exposure to lower CO concentrations may lead to the symptoms and signs of influenza or food poisoning, exposure to high concentrations of CO leads to collapse and death within minutes.

Apparently classic cases of food poisoning of a whole family may in be produced by CO poisoning. Prolonged exposure to concentrations that produce only minor symptoms may, in some cases, be associated with lasting neurological effects. These include difficulties in concentrating and emotional labiality. Complaints about such problems should alert the doctor to the possibility of CO poisoning.

Clues to the diagnosis
The following are suggestive of domestic CO poisoning:

- More than one person in the house affected;
- Symptoms disappear when away from the house, e.g. on holiday, or at work but recur on returning home;
- Symptoms related to cooking: stove in use; and

INVESTOR IN PEOPLE
POSITIVE ABOUT DISABLED PEOPLE

- Symptoms worse in winter: heating in use.

The following signs may be recognised in the home:

- Black sooty staining on or around an appliance (e.g. a stoves, boilers or fires), such as on the walls;
- Smoke or excessive condensation accumulating in rooms due to faulty flues: though you cannot smell CO, you can often smell other combustion products;
- Yellow or orange, instead of blue, flames from gas appliances or boiler pilot lights.

Clinical signs

Neurological examination is key in determining a chronic poisoning event and signs must be looked for. A neurological examination, including tests of fine movement and balance (finger–nose movement, Romberg's test, normal gait and heel–toe walking), a mini-mental state examination and testing of short-term memory and the ability to subtract 7, serially, from 100 are vital.

The cherry red skin colour is not a common sign of poisoning. This is produced when carboxyhaemoglobin (COHb) concentrations exceed about 20% and is rarely seen in life (further information in **Annex B**).

Investigations

Carbon monoxide (CO) can be measured in expired air. Breath analysers are used in smoking cessation clinics and should be used in surgeries which have such devices. There are also analysers which are available that convert CO concentration into COHb concentration from the standard equilibration curve. If such devices are used, they must be used quickly: there is no point in taking a measurement if the patient has spent hours away from the source of CO. Measurements taken the next day at the surgery may be misleading.

COHb can be measured in blood by any clinical chemistry laboratory. Venous blood should be taken into anti-coagulant and sent to the laboratory. COHb should be measured directly: measuring PO_2 (pressure of oxygen in the blood) and calculating the % saturation of haemoglobin with oxygen will be misleading as the PO_2 in CO poisoning may well be normal. Several suitable instruments are available, for example: the radiometer co-oximeter.

Pulse-oximetry in cases of suspected CO poisoning is not recommended because false high oxygen saturations are likely to be recorded due to the similar light absorbance of carboxyhaemoglobin and oxyhaemoglobin.

- Rapid measurement of expired air is useful in diagnosis.
- Blood COHb is also useful.
- Expired air CO and blood COHb are poor guides to prognosis and the need for hyperbaric treatment.

Management

- Remove patient and co-habitants from source of CO;
- Give 100% oxygen;

INVESTOR IN PEOPLE
POSITIVE ABOUT DISABLED PEOPLE

- A tightly fitting mask with an inflated face-seal is necessary for the administration of 100% oxygen;
- Consider referring for hyperbaric oxygen treatment;
- Arrange checking of appliances and flues and measurement of CO concentration in the house before allowing anyone back; and
- Contact social services, if necessary.

Indications for hyperbaric oxygen therapy (HBOT)
There is debate about the added value provided by hyperbaric oxygen. A COHb concentration of >20% should be an indication to consider hyperbaric oxygen and the decision should be taken on the basis of the indicators listed below:

- Loss of consciousness at any stage;
- Neurological signs other than headache;
- Myocardial ischaemia/arrhythmia diagnosed by ECG; or
- The patient is pregnant.

HBOT is also thought to be of use for extensive exposure to CO and if neurological damage is suspected, its use should be on a case-by-case basis.

Sources of carbon monoxide (CO)
Carbon monoxide (CO) is produced not just by malfunctioning or poorly flued gas appliances but by the incomplete combustion of all carbon-containing fuels: gas (domestic or bottled), coal, coke, oil, biofuel and wood. Stoves, fires and boilers, water heaters, paraffin heaters and room heaters are all potential sources. Caravans, boats and mobile homes are also at risk as they often use portable appliances which use these fuels and exhaust gases from vehicle engines and generators of electricity can also contain high levels of CO. During incomplete combustion, carbon, hydrogen and available oxygen combine to form carbon dioxide, water, heat and CO. Any disruption of the burning process or shortage of oxygen can increase CO production and its accumulation to dangerous levels.

Inadequate installation or maintenance of fossil fuel and wood-burning appliances leading to poor combustion of fuel, inadequate removal of waste products because of blocked and partially blocked flues and chimneys, and insufficient ventilation are the main causes of CO poisoning. Such faults can occur in all types of property, and the idea that CO poisoning is restricted to poorer homes and student accommodation is false. Owner-occupied houses with newly installed oil-powered cooking ranges can also be the site of accidents.

CO can also seep into properties via shared flues and chimneys and people may be poisoned by CO produced by an appliance in a neighbouring property. Dangerous errors, such as the venting of gas fires into cavity walls, can lead to poisoning of people living above those using the fire. Integral garages can also be a source of CO if car engines are run without adequate ventilation.

INVESTOR IN PEOPLE
POSITIVE ABOUT DISABLED PEOPLE

People to consult
For CO measurements in the house:

- ***For gas:*** Gas Safe Registered engineer – call **0800 408 5500** (Gas Safe is the Health and Safety Executive appointed provider of the gas installer registration scheme);
- Health and Safety Executive Gas Safety Advice Line – call **0800 300 363**;
- ***For oil:*** Local OFTEC (Oil Firing Technical Association) engineer – call **0845 658 5080**;
- ***For solid fuel*****:** Local HETAS (Heating Equipment Testing and Approval Scheme) engineer – call **0845 634 5626**.

Advice on the management of poisoning
Follow advice on TOXBASE (www.toxbase.org) or refer to the National Poisons Information Service (NPIS) on **0844 892 0111** for more detailed advice on the management of CO poisoning and interpretation of blood sample results.

Leaflets and further information

1. CO Awareness charity: http://www.covictim.org/

2. Gas Safe Register (has replaced CORGI registration) guidance: http://www.gassaferegister.co.uk/learn/carbon_monoxide_kills.aspx

3. *Gas Appliances – Get them checked. Keep them safe*. Leaflet produced by the Health and Safety Executive (HSE), available by calling the HSE information line on **0845 345 0055** or at: http://www.hse.gov.uk/gas/domestic/links.htm

4. HSE has also prepared a series of short videos on gas safety, which help to highlight typical scenarios and symptoms of CO poisoning: http://www.hse.gov.uk/gas/domestic/videos.htm

5. *Carbon monoxide: You can't smell it, you can't see it and it can kill!* Leaflet for information available (2008) at: http://www.documents.hps.scot.nhs.uk/environmental/sg-co-awareness-factsheet.pdf

6. Further CO guidance available on the NHS SHOW website: http://www.scot.nhs.uk/searchResults.html?cx=003501025796087152226%3Axcougsx07yc&cof=FORID%3A11&q=carbon%20monoxide&sa=Search#1147

7. Health Protection Scotland (HPS) information on CO poisoning: http://www.documents.hps.scot.nhs.uk/ewr/pdf2011/1142.pdf and http://www.hps.scot.nhs.uk/ewr/subjectsummary.aspx?subjectid=227

8. Health Protection Agency (HPA) information on CO: http://www.hpa.org.uk/web/HPAweb&Page&HPAwebAutoListName/Page/1226908886240

INVESTOR IN PEOPLE
POSITIVE ABOUT DISABLED PEOPLE

ANNEX B

Mechanisms of action of carbon monoxide (CO)

Carbon monoxide (CO) gas enters the blood system via the lung. Inhaled CO combines with haemoglobin to form carboxyhaemoglobin (COHb). Once this reaction occurs, the capacity of haemoglobin to carry oxygen is much reduced. CO binds to haemoglobin with about 240 times the affinity of oxygen and causes a left shift in the oxyhaemoglobin dissociation curve. These effects combine to reduce oxygen delivery to the tissues.

In addition, CO is transported dissolved in plasma and binds to intracellular myoglobin and mitochondrial cytochrome enzymes. Binding to cytochrome A3 is thought to play an important part in the toxicity of this gas.

Recent studies have shown that CO may function as a local transmitter substance in the body playing a role in controlling permeability of the microvasculature, and may increase adhesion of inflammatory cells and platelets to the capillary endothelium. CO poisoning leads to leakage of fluid across cerebral capillaries and thus to cerebral oedema. In those who have been exposed to enough CO to produce unconsciousness, delayed neurological damage due to leuko-encephalopathy may occur. Damage tends to be focused on those parts of the brain lying at the boundaries of the fields supplied by two cerebral arterial systems, e.g. the basal ganglia. Neurological damage seems to be the result of free radical generation and lipid peroxidation. It is possible that the binding of CO to cytochrome A3 reduces the capacity of cells to deal with free radicals.

CO bound to haemoglobin has a half-life of about 320 minutes under normal circumstances. This can be reduced by exposing the patient to 100% oxygen: this reduces the half-life to 80 minutes; or to 100% oxygen at 2 atmospheres pressure (hyperbaric oxygen), which reduces the half-life to 23 minutes. The half-life of CO bound to mitochondrial cytochromes may well be much longer than that of carboxyhaemoglobin and hyperbaric oxygen has been suggested as being important in attacking this binding site. CO binds to fetal haemoglobin and shifts the already left-shifted fetal oxyhaemoglobin dissociation curve further to the left. The half-life of CO in the fetus is longer than that in the mother.

CO is produced continuously in the body as a by-product of haem breakdown. This leads to a normal baseline COHb concentration of about 0.5%. In pregnancy and especially in haemolytic anaemias this can rise towards 5%. Cigarette smoking leads to COHb concentrations of up to about 13% in heavy smokers.

INVESTOR IN PEOPLE
POSITIVE ABOUT DISABLED PEOPLE